M. HAWKINS 14,

OPP OFFICIAL RECORD HEALTH EFFECTS DIVISION SCIENTIFIC DATA REVIEWS



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

1 8 JUN 1991

OFFICE OF PESTICIDES AND TOXIC SUBSTANCES

MEMORANDUM

Health Effects Division (HED) Peer Review Committee SUBJECT:

Draft Document on Parathion (3rd)

FROM:

TO:

Esther Rinde, Ph.D. E.R.

Manager, HED Carcinogenicity Peer Review

Science Analysis Coordination Branch Health Effects Division (H7509C)

Addressees

Attached for your review is the draft document of the Peer Review Committee on Parathion (3rd). Please provide your comments on the draft document and return to me no later than June 25, 1991. If a reply is not received by that time, we will presume that you concur and have no comments.

Should you need a few extra days for a thorough review, please let us know that your comments are forthcoming.

ADDRESSEES

- P. Fenner-Crisp
- W. Burnam
- R. Engler
- R. Hill
- K. Baetcke
- R. Beliles
- L. Brennecke
- M. Copley
- K. Dearfield
- J. Du
- B. Fisher
- G. Ghali
- H. Pettigrew
- J. Parker
- W. Sette
- M. Van Gemert
- Y. Woo
- J. Quest
- R. Zendzian

MEMORANDUM

SUBJECT: DRAFT Peer Review of Parathion (3rd)

FROM:

Esther Rinde, Ph.D. &R 6/17/9/

Science Analysis and Coordination Branch

Health Effects Division (H7509c)

TO:

Jan Auerbach, Chief Special Review Branch

Special Review and Reregistration Division (H7508c)

The Health Effects Division Peer Review Committee met on June 12, 1991 to discuss and evaluate new data on Parathion with particular reference to its bearing on the classification of Parathion's carcinogenic potential. The Committee concluded that the new data does not affect the existing classification and Parathion remains classified as a <u>Group C (possible human carcinogen) without quantification</u>, which is to say: without the use of a low dose exrapolation model (Q1*) for quantitative risk assessment; instead a reference dose (RfD) approach will be used.

A. Individuals in Attendance:

1. <u>Peer Review Committee</u>: (Signatures indicate concurrence with the peer review unless otherwise stated.)

William L. Burnam	
Reto Engler	
Marcia Van Gemert	
Karl Baetcke	
Robert Beliles	
Lucas Brennecke	
Marion Copley	
Kerry Dearfield	
Hugh Pettigrew	
Esther Rinde	****

2.	Reviewers: (Non-committee data presentation; signat accuracy of panel report.	
	rt Zendzian ice Fisher	
3.		e to attend the discussion; ncurrence with the overall
Julio Georg Richa Jean Will Yin-	lope A. Fenner-Crisp e Du ge Ghali ard Hill Parker iam Sette Tak Woo Quest	

B. Material Reviewed:

The material available for review consisted of a DER and summary memo prepared by Dr. Zendzian; Tables and statistical analysis were provided by the Registrant (HED statistician, Ms. Fisher determined that additional analysis was not warranted); Peer Review Documents for the 1st and 2nd meetings, dated Sept. 8, 1986 and July 31, 1989, respectively. The material reviewed is attached to the file copy of this report.

C. <u>Background Information</u>:

Parathion, a special review chemical, was classified as a Group C (possible human carcinogen) without quantitative risk assessment by the Peer Review Committee (PRC) at its meetings on July 1, 1986 and on March 7, 1989. A new mouse study, submitted by the Registrant, has been reviewed and the PRC was asked to consider whether the new data have any bearing on the classification of Parathion.

PRESENTATION OF NEW DATA

Reference: Carcinogenicity study of ethyl parathion administered by dosed feed to B6C3F1 mice, J.G. Page & J.E. Heath, Southern Research Institute, Project ID A21-CRM-1, 1/21/91, MRID 418173-01.

a. Experimental Design

Parathion was administered in the diet to groups of 50 male and 50 female B6C3F1 mice at 0 (control), 60, 100 and 140 ppm for 18 months.

b. Discussion of Tumor Data

The only increases in tumor occurrence that were statistically significant were observed at 60 ppm; these were lung alveolar/bronchiolar adenomas in males and systemic malignant lymphoma in females (Tables 1 and 2 - taken from the Registrant's submission). The incidence of adenomas in male mice at 60 ppm exceeded the upper range reported for historical controls at the testing facility; the incidence of malignant lymphoma in females was within the range of historical controls (Tables 3 and 4).

A <u>misdosing</u> incident, however, <u>involving this dose (60 ppm)</u> "was reported to the Agency when detected and a <u>joint decision was made</u> to continue the study" [R. Zendzian].

Dr. Zendzian quotes the Registrant's report as follows: "Due to a technical error in diet preparation, mice in the 60 ppm dose group were fed a diet containing approximately 500 ppm ethyl parathion between Days 300 and 307 of this study. The mice in this dose group were switched to control diet (i.e., untreated) diet (sic) following detection of the misdosing incident for 17 days (Days 300 to 323) to allow these mice to regain lost weight and recover from the acute toxicity induced by exposure to an excessive dose of test article. The incident is highlighted here because it had an impact on study parameters (e.g., mortalities, body weights, food consumption) and study interpretation." (Six males and two females died within 14 days of the misdosing.)

c. <u>Consideration of Adequacy of Dosing for Assessment of</u> Carcinogenic Potential:

Both male and female mice were considered to have been adequately, but not excessively dosed (at 140 ppm), based on dose-related decreases (less than 5%) in body-weight gain and significant brain, plasma and RBC cholinesterase inhibition (without a compound induced increase in mortality).

The low dosed animals ("60" ppm) were of course overdosed.

Table 1:

CARCINOGENICITY STUDY OF ETHYL PARATHION ADMINISTERED BY DOSED FEED TO B6C3F1 MICE Study: A21-CRM-1

Analysis of Histopathology Findings - Males Tumor Pathology

	<u>Untreated</u>	60 PPM	100 PPM_	140_PPM
Lung - Alveolar/Bronchiolar Adenoma		· · · · · · · · · · · · · · · · · ·	, , , , , , , , , , , , , , , , , , , 	the second secon
Overall Rates (a) Terminal Rates (b) Life Table Test (c)	5/50 (10.0%) 5/49 (10.2%)	13/50 (26.0%) 12/39 (30.8%) p = 0.009*	6/50 (12.0%) 6/47 (12.8%) p = 0.471	4/50 (8.0%) 4/50 (8.0%) p = 0.513
Incidental Tumor Test (d) Cochran-Armitage Trend Test (e) Fisher Exact Test (e)	p = 0.206	p = 0.014* p = 0.033*	p = 0.457 p = 0.500	р = 0.500 р = 0.500
<u>Lung -</u> Alveolar/Bronchiolar Carcinoma			• • • • • • • • • • • • • • • • • • • •	. •
Overall Rates (a) Terminal Rates (b) Life Table Test (c)	0/50 0/49	1/50 (2.0%) 1/39 (2.6%) p = 0.454	0/50 0/47 p = 1.000	0/50 0/50 p = 1.000
Incidental Tumor Test (d) Cochran-Armitage Trend Test (e) Fisher Exact Test (e)	р = 0.500	p = 0.450 p = 0.500	p = 1.000 p = 1.000	p = 1.000 p = 1.000
<u>lung -</u> <u>Alveolar/Bronchiolar Adenoma and</u> <u>Carcinoma Combined</u>				
Overall Rates (a) Terminal Rates (b) Life Table Test (c)	5/50 (10.0%) 5/49 (10.2%)	14/50 (28.0%) 13/39 (33.3%) p = 0.005*	6/50 (12.0%) 6/47 (12.8%) p = 0.471	4/50 (8.0%) 4/50 (8.0%) p = 0.513
Incidental Tumor Test (d) Cochran-Armitage Trend Test (e) Fisher Exact Test (e)	p = 0.185	p = 0.007* p = 0.020*	p = 0.457 p = 0.500	р = 0.500 р = 0.500

133-A21CRM1

Table 2
CARCINOGENICITY STUDY OF ETHYL PARATHION ADMINISTERED BY DOSED FEED TO B6C3F1 MICE
Study: A21-CRM-1
Analysis of Histopathology Findings - Females
Tumor Pathology

-		<u>Untreated</u>	<u>60 PPH</u>	100 PPM	140 PPH	
	Harderian Gland - Adenoma					
-	Overall Rates (a) Terminal Rates (b) Life Table Test (c) Incidental Tumor Test (d) Cochran-Armitage Trend Test (e) Fisher Exact Test (e)	1/50 (2.0%) 1/44 (2.3%) p = 0.170	1/50 (2.0%) 1/42 (2.4%) p = 0.751 p = 0.756 p = 0.753	0/50 0/48 p = 0.517 p = 0.517	0/50 0/44 p = 0.500 p = 0.500 ρ = 0.500	145
⇒	<u>Systemic -</u> <u>Lymphoma Halignant</u>				en e	5 145-A21CRMI
-	Overall Rates (a) Terminal Rates (b) Life Table Test (c) Incidental Tumor Test (d) Cochran-Armitage Trend Test (e) Fisher Exact Test (e)	0/50 0/44 p - 0.332	5/50 (10.0%) 3/42 (7.1%) p = 0.033* p = 0.033* p = 0.028*	3/50 (6.0%) 2/48 (4.2%) p = 0.140 p = 0.060 p = 0.121	2/50 (4.0%) 1/44 (2.3%) p = 0.242 p = 0.240 p = 0.247	ā
	<u>Systemic -</u> <u>Histiocytic Sarcoma</u>	i		Commence Spiritual Commence Co	. gas at a fee and decide a con-	
	Overall Rates (a) Terminal Rates (b) Life Table Test (c) Incidental Tumor Test (d)	0/50 0/44	1/50 (2.0%) 0/42 p = 0.504 p = 0.500	0/50 0/48 p = 1.000 p = 1.000	2/50 (4.0%) 0/44 p = 0.249 p = 0.223	
	Cochran-Armitage Trend Test (e) Fisher Exact Test (e)	p = 0.149	ę = 0.500	p - 1.000	p - 0.247	

VERSION: 09/11/50	TOXICOLOGY DATA MANAGEMENT SYSTEM TUMOR INCIDENCE FOR BELECTED CONTROL MINAL GROUPS	PAGE: 441
CONTRACT/LAS: SRI SPECIES: MICE STRAIN: MICE-BECOFI	ROUTE! ORAL Vehicle: Feed	DATE OF INPUT FILE: 15-5EP-1990
LENGTH OF STUDY, CHRONIC	INCIDENCE DATA FOR VEHICLE CONTROL GROUPS	
	MTX	FEHALE
LIVER: MEPATOCELLULAR CARCINOMA, B HEPATOCELLULAR ADENOMA, HEPATOCELLULAR ADENOMA, HEPATOCELLULAR ADENOMA, HERE AD	10/50 (201) 13/49 (271) 12/50 (241)	2/50 (41) 6/50 (161) 3/50 (61) [
TOTAL & INCIDENCE	35/149 (234) B.D 3.51	13/150 (91) B.D 6.43
*Liver; Bepatocholangiocalcinoma Total Incidence	0/80 (01) 0/49 (01) 0/50 (01) 0/149 (01) 8.D 0.00	0/50 (01) 0/50 (01) 0/30 (01) 0/150 (01) 8.D 0.00
"LIVER: ITO CELL TUMOR BENIGH TOTAL \ INCIDENCE	0/50 (01) 0/49 (01) 0/50 (01) 0/149 (01) 8.D 0.00	0/50 (01) 0/50 (01) 0/50 (01) 1 0/150 (01) 8.D 0.00
*LIVER; LEVENIA ENTHROCITIC TOTAL ^ INCIDENCE	0/50 (01) 0/49 (01) 0/50 (01) 0/149 (01) 8.D. = 0.00	0/50 (01) 0/50 (01) 0/50 (01) 0/150 (01) 8.D 0.00
"LIVER; SANCOMA TOTAL & INCIDENCE	0/50 (01) 0/49 (01) 0/50 (01) 0/149 (01) 8.D. = 0.00	0/50 (01) 0/50 (01) 0/50 (01) 0/150 (01) 8.D 0.00
*LUNG! ALVEGLAR/BRONCETOLAR ADRIONA TOTAL & INCIDENCE	8/50 (101) 6/50 (121) 5/50 (101) 16/150 (111) 8.0 1.15	1/50 (21)
*LUNG: ALVECOLA SECONCISTOLAR ADGROSA OR ADERONA TOTAL & INCIDENCE	8/50 (10t) 6/50 (12t) 5/50 (10t) 16/150 (11t) 8.D 1.15	2/50 (41) 3/50 (61) 1/50 (21) Y
*LUBG! ALVEGIAR/BRONCHIOLAR CARCÍNOHA TOTAL & INCIDENCE	3/80	2/50 (40)
TUMG ALVEGIAN/BROMCHIOLAR ALVEGIANGA OR ALVEGIAR/BROMCHIOLAR	6/50 (124) 9/50 (184) 11/50 (224)	3/50 (61) 4/50 (61) 3/50 (61)
TOTAL & INCIDENCE	26/150 (174) 6.0 6.03	10/150 (74) 6.D 1.15
CANCINONA CANCINONA TOTAL 1 INCIDENCE	0/50 (01) 0/50 (01) 0/50 (01) 0/150 (01) 8.D 0.00	0/50 (01) 0/50 (01) 0/50 (01) 0/150 (01) 8.D. = 0.00
LUNG: CARCINOMA, ALVEOLAR/BRONCHIOLAR CARCINOMA OR ADENOSQUANOUS	1/50 (21) 3/50 (61) 6/50 (121)	1/50 (21) 1/50 (21) 2/50 (41)
CANCINGAIA TOTAL & INCIDENCE	10/150 (71) 4,D, - 5,03	4/150 (31) 8,D, - 1,15

4. DENOMINATOR IS NUMBER OF ANIMALS WITH TISSUES EXAMINED MICROSCOPICALLY 1. DENOMINATOR IS NUMBER OF ANIMALS NECROPSIED

		_	_	_	_	_	_		_
1	16	٥.	Δ	7	1	r	D	М	1

	VERSION: 09/17/90	TOKON INCIDENCE FOR SELECTED CONTROL ANIMAL GROUPS	ACEMENT SYSTEM D CONTROL ANIHAL (sanous				2	PAGE: 433	
	CONTRACT/LAB: SRI SPECIES: MICE STARIN: HICE-B6CSFI	MOUTE: DRAL. VENICLE: FEED	9			DATE OF INPUT F	DATE OF INPUT FILE: 15-8EP-1990	15-81	CP-1990	
	LENGTH OF STUDY: CHRONIC	INCIDENCE DATA FOR VEHICLE CONTROL GROUPS	CLE CONTROL GROUP							
		HAYK				PEHALE				_
	HORS CIDENCE	12/50 (241) 18/50 (341) 1 46/150 (321) 8,D, = 6	10/50 (361)	15/50 ((301) 13/5/ 39/150 (261)			11/50	(221)	
	TALL CRIMINT TUMORS TOTAL & INCIDENCE		26/50 (521) 4.00	21/50 (421)		21/50 (421) (401) 8.D.		18/50 (361) 3,46	(361)	
	TOTAL CHOINE TALLOHANT AND REVIOU TUVORA TALLOHANT AND REVIOU TUVORAL	31/80 (621) 39/80 (781) 3 305/150 (701) 8.D 8	35/50 (701) 8.00	30/50 ((401)	29/50 (1	(301) 8.0	24/50	(487)	
		0/50 (01) 1/50 (21) 1/150 (11) 8,B, - 1	6/50 (01) 1.15	1/30 ((21) 1/50 (21) 2/150 (11) 8,6	1/50 (2	21) 8.b. =	0/50 1.15	(00)	
		3/50 (61) 1/50 (21) 7/150 (51) 8.D. = 2	3/50 (61) 2,31	2/50 (2/50 (41) 2/50 4/150 (31)	2/50 (4	(41) 8.b, -	0/50 2,31	(01)	<u>, </u>
	TAXL GRANE: Hemanologracom or remanolom Total, 1 incluence		3/50 (61) 1,15	3/50 (61) 5/150	61) (150 (3	2/50 (((44) 5.0. =	0/30 3.06	(00)	
		0/50 (01) 0/30 (01) 0/150 (01) 2,D, = 0	0/50 (01)	0/50 (01)	01) /150 (0	_	(01) E.D. =	0/50 0.00	(10)	
		1/50 (21) 0/10 (01)	0/20 (01)	0/30 (01)		0/20 (0/)	ž	(10) 05/0	(0)	
	OR UNDIFFERENTIATED TOTAL & INCIDENCE	1/150 (11) 6.9 1.	1.19		(10) 051/0	ł	6.0. =	9.0	,	
A	TALL ONCANS NATIONAL AND NISTICOTIS SAUCOMA NISTICOTIS SAUCOMA OTAL	5/50 (101) 5/50 (101) 15/150 (101) 5.D 0	5/56 (10t) 6.00	12/50 (241)	_		1	13/50 (261)	(261)	\bigvee
ے۔	TALL ORGANS: MALIGHANT LIMPHONA: MASTOCTIC, LIMPHOCITIC, MISTO, OR UNDIFFERNITATED	ă	1101) 95/9	12/50 (241)		16/50 (321)		13/50 (261)	(36/)	i
	CELL TYPE	15/150 (101) 8.D 0	00.0	•	41/150 (274) 8.D	171	- i.	4.16		
-	HALL ORGINS: HESOTHELIONA PENTON, MALICHANT, NOS TOTAL & INCIDENCE	0/56 (01) 0/30 (01) 0/150 (01) 8.0. = 0	0/50 (01) 0,00	0/20 ((01) 0/50 (01) 0/150 (01) 8.D.	0/20 (0	 	0,00	(10)	

* DENOMINATOR IS NUMBER OF ANIMALS WITH TISSUES EXAMINED MICROSCOPICALLY IS DENOMINATOR IS NUMBER OF ANIMALS MECHOFSIED

Conclusion

The PRC concluded that, despite the misdosing, the new mouse study was acceptable since, the incident was reported to the Agency at the point of discovery by the Registrant, at which time the Agency agreed that the study could be continued.

Although the tumors occurring at the low dose were considered to be difficult to interpret, nevertheless, it was agreed that they could not be discounted. The PRC unanimously concluded that this new information does not affect the existing classification of Parathion as <u>Group C (possible human carcinogen)</u>, <u>without quantification</u>, which is to say: without the use of a low dose exrapolation model (Q1*) for quantitative risk assessment; instead a reference dose (RfD) approach will be used.

Previously submitted data, and its contribution to the weight of evidence for Parathion, were fully discussed in the memos documenting Peer Review Meetings 1 and 2 (dated Sept. 8, 1986 and July 31, 1989, respectively).



012431

Chemical:

Parathion

PC Code:

HED File Code

Memo Date:

File ID:

Accession Number:

057501

21200 PEER REVIEW

06/18/1991

00000000

412-01-0123

HED Records Reference Center 03/20/2001